

Efficient symmetric multiparty quantum state sharing of an arbitrary m -qubit state

Xi-Han Li,^{1,2} Ping Zhou,^{1,2} Chun-Yan Li,^{1,2} Hong-Yu Zhou,^{1,2,3} and Fu-Guo Deng^{1,2,3*}

¹ *The Key Laboratory of Beam Technology and Material Modification of Ministry of Education, Beijing Normal University, Beijing 100875, People's Republic of China*

² *Institute of Low Energy Nuclear Physics, and Department of Material Science and Engineering, Beijing Normal University, Beijing 100875, People's Republic of China*

³ *Beijing Radiation Center, Beijing 100875, People's Republic of China*

(Dated: February 1, 2008)

We present a scheme for symmetric multiparty quantum state sharing of an arbitrary m -qubit state with m Greenberger-Horne-Zeilinger states following some ideas from the controlled teleportation [Phys. Rev. A **72**, 02338 (2005)]. The sender Alice performs m Bell-state measurements on her $2m$ particles and the controllers need only to take some single-photon product measurements on their photons independently, not multipartite-entanglement measurements, which makes this scheme more convenient than the latter. Also it does not require the parties to perform a controlled-NOT gate on the photons for reconstructing the unknown m -qubit state and it is an optimal one as its efficiency for qubits approaches 100% in principle.

PACS numbers: 03.67.Hk, 03.67.Dd, 03.65.Ud, 89.70.+c

I. INTRODUCTION

Suppose a president of a bank, say Alice wants to send some secret message M_A to her agents, Bob and Charlie who are at a remote place for her business. She doubts that there may be one of the two persons dishonest and he will destroy the business with the message independently. Alice believes that the honest one will keep the dishonest one from doing any damage if they both appear in the process of dealing with the business. In this way, the message cannot be transmitted to the agents directly. In classical secret sharing [1], Alice splits M_A into two pieces, M_B and M_C , and sends them to Bob and Charlie, respectively. When they act in concert, they can recover the message $M_A = M_B \oplus M_C$; otherwise, they can obtain nothing about the message. The classical signal is in one of the eigenvectors of a measuring basis (MB), say σ_z , and it can be copied fully and freely [2]. It is impossible for the parties to communicate in an unconditionally secure way if they only resort to classical communication [3]. When quantum mechanics enters the field of information, the story is changed [3, 4].

Quantum secret sharing (QSS) is the generalization of classical secret sharing into quantum scenario. Most existing QSS schemes are focused on creating a private key among several parties or splitting a classical secret. For example, an original QSS scheme [5] was proposed by Hillery, Bužek, and Berthiaume (HBB) in 1999 by using three-particle and four-particle entangled Greenberger-Horne-Zeilinger (GHZ) states for distributing a private key among some agents and sharing classical information. In HBB scheme, the three parties, Alice, Bob and Charlie choose randomly two measuring bases (MBs), σ_x and σ_y to measure their particles independently. The probabil-

ity that the quantum resource can be used for carrying the useful information is 50%. That is, their results are correlated and will be kept for creating a key when they all choose the MB σ_x or one chooses σ_x and the others choose σ_y , which takes place with the probability 50%. They removed the ideas in the controlled teleportation [6] to split a classical secret among the agents. Subsequently, Karlsson, Koashi and Imoto proposed another QSS protocol for those two goals with multi-particle entangled states and entanglement swapping [7]. Its intrinsic efficiency for qubits η_q [8], the ratio of number of theoretical valid transmitted qubits to the number of transmitted qubits is about 50% as half of the instances will be abandoned [9]. Now, there are some theoretic schemes for sharing and splitting a classical message [5, 7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22]. The experiment demonstration of QSS was also studied by some groups [23, 24].

Recently, a novel concept, quantum state sharing (QSTS) was proposed and actively pursued by some groups [24, 25, 26, 27]. It is the extension of QSS for sharing an unknown state among several agents by resorting to the multipartite entanglements. Cleve, Gottesman and Lo [25] introduced a way for a (k, n) threshold QSTS scheme which can be used to split a secret quantum state related to a classical secret message [27]. Li et al. [26] proposed a scheme for sharing an unknown single qubit with some Einstein-Podolsky-Rosen (EPR) pairs and a multi-particle joint measurement. In 2004, Lance et al. studied QSTS with continuous variable [24]. Deng et al. [27] introduced a scheme for splitting an arbitrary two-qubit state with EPR pairs and GHZ-state measurements. Also, some controlled teleportation schemes were discussed in Ref.[6, 28, 29]. The unknown state can be teleported to the sender with the control of some controllers. In fact, almost all those controlled teleportation schemes can be used for QSTS with or without a little modification. For example, in Ref. [29], QSTS for an

*Email: fgdeng@bnu.edu.cn

arbitrary two-qubit state can be implemented with the symmetric controlled teleportation scheme [29] without any modification. In essence, a secure QSTS scheme can also be used for controlled teleportation by means that some agents act as the controllers in the latter.

In this paper, we will present a scheme for QSTS of an arbitrary m -particle state following some ideas in Ref. [29]. It will be shown that the agents need only to perform a product measurement $\sigma_x^1 \otimes \sigma_x^2 \otimes \dots \otimes \sigma_x^m$ on their particles, not multipartite-entanglement measurements, which makes this scheme more convenient than Ref. [29]. Moreover, it requires the agents only to perform some local unitary operations on the photons retained for recovering the unknown state, not a two-qubit joint operation, such as controlled-NOT operation.

II. QSTS OF AN ARBITRARY M -QUBIT STATE

A. QSTS of an arbitrary two-qubit state with two agents

For simplicity, we first present a way for the symmetric quantum state sharing of an arbitrary two-qubit state with two three-particle GHZ states. That is, there are two agents, Bob and Charlie. We assume that the parties share a sequence of multipartite entangled states securely first, similarly as Refs. [29, 30]. Similar to quantum secure direct communication [30, 31], the parties can do quantum privacy amplification on the sequence of entangled states shared with quantum entanglement purification [32, 33] in a noise channel.

The basic idea of this QSTS scheme can be described as following. Suppose that the quantum information is an unknown arbitrary two-qubit state

$$|\Phi\rangle_{xy} = a|00\rangle_{xy} + b|01\rangle_{xy} + c|10\rangle_{xy} + d|11\rangle_{xy}, \quad (1)$$

where $|0\rangle$ and $|1\rangle$ are the two eigenvector of the MB σ_z (for example the polarization of the single photon along the z -direction), and

$$|a|^2 + |b|^2 + |c|^2 + |d|^2 = 1. \quad (2)$$

Alice prepares two three-particle GHZ states

$$|\Psi\rangle_{a_1 a_2 a_3} = |\Psi\rangle_{b_1 b_2 b_3} = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle). \quad (3)$$

Same as Ref. [29], Alice sends the photons a_2 and b_2 to Bob, and the photons a_3 and b_3 to Charlie, respectively. She performs Bell-state measurements on the photons x and a_1 , and y and b_1 , then the state of the unknown quantum system $|\Phi\rangle_{xy}$ is transferred to the quantum system composed of the four photons a_2 , a_3 , b_2 , and b_3 . When the two agents want to recover the unknown state $|\Phi\rangle_{xy}$, one of them performs two single-photon measurements on his photons and the other can get the state with two local unitary operations. In detail, the state of the composite quantum system composed of the eight photons $x, y, a_1, a_2, a_3, b_1, b_2$ and b_3 can be written as

$$\begin{aligned} |\Psi\rangle_s &\equiv |\Phi\rangle_{xy} \otimes |\Psi\rangle_{a_1 a_2 a_3} \otimes |\Psi\rangle_{b_1 b_2 b_3} \\ &= (a|00\rangle + b|01\rangle + c|10\rangle + d|11\rangle)_{xy} \\ &\quad \otimes \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)_{a_1 a_2 a_3} \\ &\quad \otimes \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)_{b_1 b_2 b_3}. \end{aligned} \quad (4)$$

When Alice performs the Bell-state measurement on the photons x and a_1 , they are randomly in one of the four Bell states

$$|\phi^\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle|0\rangle \pm |1\rangle|1\rangle), \quad (5)$$

$$|\psi^\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle|1\rangle \pm |1\rangle|0\rangle). \quad (6)$$

So does the quantum system composed of the photons y and b_1 .

As an example for demonstrating the principle of this QSTS scheme, we assume that the outcomes obtained by Alice are $|\phi^+\rangle_{xa_1}$ and $|\phi^+\rangle_{yb_1}$. Then the retained four photons are in the state $|\Psi\rangle_r$. Here

$$\begin{aligned} |\Psi\rangle_r &= a|00\rangle_{a_2 b_2}|00\rangle_{a_3 b_3} + b|01\rangle_{a_2 b_2}|01\rangle_{a_3 b_3} \\ &\quad + c|10\rangle_{a_2 b_2}|10\rangle_{a_3 b_3} + d|11\rangle_{a_2 b_2}|11\rangle_{a_3 b_3}. \end{aligned} \quad (7)$$

If Bob performs the product measurement $\sigma_x \otimes \sigma_x$ on the photons a_2 and b_2 , the state $|\Psi\rangle_r$ can be written as

$$\begin{aligned} |\Psi\rangle_r &= \frac{1}{2} [|+\rangle_{a_2}|+\rangle_{b_2}(a|00\rangle_{a_3 b_3} + b|01\rangle_{a_3 b_3} + c|10\rangle_{a_3 b_3} + d|11\rangle_{a_3 b_3}) \\ &\quad + |+\rangle_{a_2}|-\rangle_{b_2}(a|00\rangle_{a_3 b_3} - b|01\rangle_{a_3 b_3} + c|10\rangle_{a_3 b_3} - d|11\rangle_{a_3 b_3}) \\ &\quad + |-\rangle_{a_2}|+\rangle_{b_2}(a|00\rangle_{a_3 b_3} + b|01\rangle_{a_3 b_3} - c|10\rangle_{a_3 b_3} - d|11\rangle_{a_3 b_3}) \\ &\quad + |-\rangle_{a_2}|-\rangle_{b_2}(a|00\rangle_{a_3 b_3} - b|01\rangle_{a_3 b_3} - c|10\rangle_{a_3 b_3} + d|11\rangle_{a_3 b_3})]. \end{aligned} \quad (8)$$

That is, Charlie can recover the unknown state $|\Phi\rangle_{xy}$

with two local unitary operations if Bob and Charlie act

TABLE I: The relation between the local unitary operations and the results R_{xa_1} , R_{yb_1} , R_{a_2} and R_{b_2} . $\Phi_{a_3b_3}$ is the state of the two particles hold in the hand of Charlie after all the measurements are done by Alice and Bob; U_C are the local unitary operations with which Charlie can reconstruct the unknown state $|\Phi\rangle_{xy}$.

V_{xa_1}	V_{yb_1}	$P_{xa_1} \otimes P_{a_2}$	$P_{yb_1} \otimes P_{b_2}$	$\Phi_{a_3b_3}$	U_C
0	0	+	+	$a 00\rangle + b 01\rangle + c 10\rangle + d 11\rangle$	$U_0 \otimes U_0$
0	0	+	-	$a 00\rangle - b 01\rangle + c 10\rangle - d 11\rangle$	$U_0 \otimes U_1$
0	0	-	+	$a 00\rangle + b 01\rangle - c 10\rangle - d 11\rangle$	$U_1 \otimes U_0$
0	0	-	-	$a 00\rangle - b 01\rangle - c 10\rangle + d 11\rangle$	$U_1 \otimes U_1$
0	1	+	+	$a 01\rangle + b 00\rangle + c 11\rangle + d 10\rangle$	$U_0 \otimes U_2$
0	1	+	-	$a 01\rangle - b 00\rangle + c 11\rangle - d 10\rangle$	$U_0 \otimes U_3$
0	1	-	+	$a 01\rangle + b 00\rangle - c 11\rangle - d 10\rangle$	$U_1 \otimes U_2$
0	1	-	-	$a 01\rangle - b 00\rangle - c 11\rangle + d 10\rangle$	$U_1 \otimes U_3$
1	0	+	+	$a 10\rangle + b 11\rangle + c 00\rangle + d 01\rangle$	$U_2 \otimes U_0$
1	0	+	-	$a 10\rangle - b 11\rangle + c 00\rangle - d 01\rangle$	$U_2 \otimes U_1$
1	0	-	+	$a 10\rangle + b 11\rangle - c 00\rangle - d 01\rangle$	$U_3 \otimes U_0$
1	0	-	-	$a 10\rangle - b 11\rangle - c 00\rangle + d 01\rangle$	$U_3 \otimes U_1$
1	1	+	+	$a 11\rangle + b 10\rangle + c 01\rangle + d 00\rangle$	$U_2 \otimes U_2$
1	1	+	-	$a 11\rangle - b 10\rangle + c 01\rangle - d 00\rangle$	$U_2 \otimes U_3$
1	1	-	+	$a 11\rangle + b 10\rangle - c 01\rangle - d 00\rangle$	$U_3 \otimes U_2$
1	1	-	-	$a 11\rangle - b 10\rangle - c 01\rangle + d 00\rangle$	$U_3 \otimes U_3$

in concert. When the outcomes of the product measurement $\sigma_x \otimes \sigma_x$ done by Bob are $|+x\rangle|+x\rangle$, $|+x\rangle|-x\rangle$, $|-x\rangle|+x\rangle$, and $|-x\rangle|-x\rangle$, Charlie should perform the unitary operations $U_0 \otimes U_0$, $U_0 \otimes U_1$, $U_1 \otimes U_0$, and $U_1 \otimes U_1$, respectively on the photons a_3 and b_3 to transfer them to the unknown quantum system. Here

$$U_0 = |0\rangle\langle 0| + |1\rangle\langle 1|, \quad U_1 = |0\rangle\langle 0| - |1\rangle\langle 1|. \quad (9)$$

For the other case, the relation between the measurement results and the final state and the operations needed to reconstruct the original state $|\Phi\rangle_{xy}$ is shown in Table I. Similar to Ref. [29], we define V as the bit value of the Bell state, i.e., $V_{|\phi^\pm\rangle} \equiv 0$, $V_{|\psi^\pm\rangle} \equiv 1$. That is, the bit value $V = 0$ if the states of the two particles are parallel, otherwise $V = 1$. P denotes the parity of the Bell-basis measurement and the single-particle measurement σ_x , i.e.,

$$P_{|\phi^\pm\rangle} \equiv \pm, \quad P_{|\psi^\pm\rangle} \equiv \pm, \quad P_{|\pm x\rangle} \equiv \pm, \quad (10)$$

and

$$U_2 = |1\rangle\langle 0| + |0\rangle\langle 1|, \quad U_3 = |0\rangle\langle 1| - |1\rangle\langle 0|. \quad (11)$$

Different from Ref. [29], it is unnecessary for Alice to take a Hadamard operation on each photon in the GHZ state in this QSTS scheme. Also, the agents need not take a controlled-NOT (CNOT) gate on the two photons retained for transferring them to the originally unknown two-qubit state $|\Phi\rangle_{xy}$, only two local unitary operations, when the parties cooperate. Moreover, the agents need only two single-photon measurements, not multipartite entanglement measurements, which makes this QSTS scheme more convenient than that in Ref. [29]. Also, its efficiency for qubits η_q is 100% as all the quantum resources are useful in theory.

B. QSTS of an arbitrary two-qubit state with $n+1$ agents

It is straightforward to generalize this QSTS scheme to the case with $n+1$ agents, similar to Ref. [29]. For this multi-party QSTS, Alice will prepare two $(n+2)$ -photon GHZ states and share them with the $n+1$ agents securely

first. Then the composite state of the system is

$$\begin{aligned} |\Psi\rangle_S &\equiv |\Phi\rangle_{xy} \otimes |\Psi\rangle_{s_1} \otimes |\Psi\rangle_{s_2} \\ &= (a|00\rangle + b|01\rangle + c|10\rangle + d|11\rangle)_{xy} \\ &\quad \otimes \frac{1}{\sqrt{2}} \left(\prod_{i=1}^{n+2} |0\rangle_{a_i} + \prod_{i=1}^{n+2} |1\rangle_{a_i} \right) \end{aligned}$$

$$\otimes \frac{1}{\sqrt{2}} \left(\prod_{i=1}^{n+2} |0\rangle_{b_i} + \prod_{i=1}^{n+2} |1\rangle_{b_i} \right) \quad (12)$$

After Alice takes the Bell-state measurements on the particles x and a_1 , and y and b_1 , respectively, the state of the subsystem (without being normalized) becomes

$$\begin{aligned} \Psi_{sub} = & \alpha \prod_{i=2}^{n+2} |0\rangle_{a_i} \prod_{i=2}^{n+2} |0\rangle_{b_i} + \beta \prod_{i=2}^{n+2} |0\rangle_{a_i} \prod_{i=2}^{n+2} |1\rangle_{b_i} \\ & + \gamma \prod_{i=2}^{n+2} |1\rangle_{a_i} \prod_{i=2}^{n+2} |0\rangle_{b_i} + \delta \prod_{i=2}^{n+2} |1\rangle_{a_i} \prod_{i=2}^{n+2} |1\rangle_{b_i} \end{aligned} \quad (13)$$

The relation between the parameters $\alpha, \beta, \gamma, \delta$ and the results R_{xa_1} and R_{yb_1} is shown in Table II.

When the agents want to reconstruct the original state $|\Phi\rangle_{xy}$, n agents perform the single-photon product measurement $\sigma_x \otimes \sigma_x$ on their photons independently.

Let us assume that the agents who measure the photons are Bob _{i} ($i = 2, 3, \dots, n+1$) and the one who does not measure his photons is Charlie. That is, the Bob _{i} act as the controllers for recovering the unknown state. The measurements done by them can be express by the formula M

$$M \equiv [(\langle +x|)^{n-t} (\langle -x|)^t]_a \otimes [(\langle +x|)^{n-q} (\langle -x|)^q]_b. \quad (14)$$

Here $[(\langle +x|)^{n-t} (\langle -x|)^t]_a$ is the measurement operation related to the state of a_i , and $[(\langle +x|)^{n-q} (\langle -x|)^q]_b$ is related to b_i . t and q are the numbers of the controllers who obtain the result $\langle -x|$ when they measure the particle a_i and b_i , respectively. After the measurements done by Alice and the n controllers, the final state $\Phi_{a_{n+2}b_{n+2}}$ can be gained by means of performing the operation M on the state Ψ_{sub} ,

$$\begin{aligned} \Phi_{a_{n+2}b_{n+2}} = & M \left(\alpha \prod_{i=2}^{n+2} |0\rangle_{a_i} \prod_{i=2}^{n+2} |0\rangle_{b_i} + \beta \prod_{i=2}^{n+2} |0\rangle_{a_i} \prod_{i=2}^{n+2} |1\rangle_{b_i} + \gamma \prod_{i=2}^{n+2} |1\rangle_{a_i} \prod_{i=2}^{n+2} |0\rangle_{b_i} + \delta \prod_{i=2}^{n+2} |1\rangle_{a_i} \prod_{i=2}^{n+2} |1\rangle_{b_i} \right) \\ = & \alpha |0\rangle_{a_{n+2}} |0\rangle_{b_{n+2}} + (-1)^q \beta |0\rangle_{a_{n+2}} |1\rangle_{b_{n+2}} + (-1)^t \gamma |1\rangle_{a_{n+2}} |0\rangle_{b_{n+2}} + (-1)^{q+t} \delta |1\rangle_{a_{n+2}} |1\rangle_{b_{n+2}}. \end{aligned} \quad (15)$$

We define

$$P_1 = P_{xa_1} \otimes \prod_{i=2}^{n+1} P_{a_i}, \quad P_2 = P_{yb_1} \otimes \prod_{i=2}^{n+1} P_{b_i}, \quad (16)$$

where P_{xa_1} and P_{yb_1} are the results of Bell-basis measurements done by Alice, and P_{a_i} and P_{b_i} are the results of single-particle measurements done by the n controllers, respectively.

The relation between the results of the measurements and the local operations with which Charlie reconstructs the original state when he cooperate with Bob _{i} is as the same as that in Table I with a little modification. That is, $P_{xa_1} \otimes P_{a_2}$, $P_{yb_1} \otimes P_{b_2}$, and $\Phi_{a_3b_3}$ are replaced by P_1 , P_2 , and $\Phi_{a_{n+2}b_{n+2}}$, respectively.

Same as the case with two agents, this multiparty QSTS does not require the agents to do Bell-state measurements on their photons. Whether the number of the controllers is odd or even, the last agent Charlie need only take two local unitary operations on the two photons for reconstructing the unknown two-qubit state.

TABLE II: The relation between the values of $\alpha, \beta, \gamma, \delta$ and the results of Bell-basis measurement R_{xa_1} and R_{yb_1}

V_{xa_1}	V_{yb_1}	P_{xa_1}	P_{yb_1}	α	β	γ	δ
0	0	+	+	$+a$	$+b$	$+c$	$+d$
0	0	+	-	$+a$	$-b$	$+c$	$-d$
0	0	-	+	$+a$	$+b$	$-c$	$-d$
0	0	-	-	$+a$	$-b$	$-c$	$+d$
0	1	+	+	$+b$	$+a$	$+d$	$+c$
0	1	+	-	$-b$	$+a$	$-d$	$+c$
0	1	-	+	$+b$	$+a$	$-d$	$-c$
0	1	-	-	$-b$	$+a$	$+d$	$-c$
1	0	+	+	$+c$	$+d$	$+a$	$+b$
1	0	+	-	$+c$	$-d$	$+a$	$-b$
1	0	-	+	$-c$	$-d$	$+a$	$+b$
1	0	-	-	$-c$	$+d$	$+a$	$-b$
1	1	+	+	$+d$	$+c$	$+b$	$+a$
1	1	+	-	$-d$	$+c$	$-b$	$+a$
1	1	-	+	$-d$	$-c$	$+b$	$+a$
1	1	-	-	$+d$	$-c$	$-b$	$+a$

C. QSTS of an arbitrary m -qubit state with $n + 1$ agents

An m -qubit state can be described as

$$|\Phi\rangle_u = \sum_{ij\dots k} a_{ij\dots k} \underbrace{|ij\dots k\rangle}_m |ij\dots k\rangle_{x_1 x_2 \dots x_m}, \quad (17)$$

where $i, j, \dots, k \in \{0, 1\}$, and $x_1, x_2, \dots$, and x_m are the m particles in the unknown state. For sharing this m -qubit state, Alice will prepare m $(n + 2)$ -photon GHZ states and share them with the $n + 1$ agents, say Bob $_j$ ($j = 1, 2, \dots, n$) and Charlie. Then the state of the composite quantum system can be written as

$$\begin{aligned} |\Psi\rangle \equiv & \left(\sum_{ij\dots k} a_{ij\dots k} \underbrace{|ij\dots k\rangle}_m |ij\dots k\rangle_{x_1 x_2 \dots x_m} \right) \\ & \otimes \prod_{i=1}^m \left[\frac{1}{\sqrt{2}} \left(\prod_{j=0}^{n+1} |0\rangle_{b_{ij}} + \prod_{j=0}^{n+1} |1\rangle_{b_{ij}} \right) \right], \end{aligned} \quad (18)$$

where the particles b_{ij} ($i = 1, 2, \dots, m$) are held in the hand of Bob $_j$ ($j = 1, 2, \dots, n$), b_{i0} are the particles of Alice's, and b_{in+1} are controlled by Charlie. Similar to Ref. [34], after the Bell-state measurements on the particles x_i and b_{i0} done by Alice ($i = 1, 2, \dots, m$), the unknown m -qubit state will be transferred to the quantum system composed of the photons kept by the agents. Each of Bob $_j$ performs the product measurement $\sigma_x^1 \otimes \sigma_x^2 \otimes \dots \otimes \sigma_x^m$ on his m photons b_{ij} ($i = 1, 2, \dots, m$), then the state of the photons kept by Charlie becomes $(U_1'^{-1} \otimes \dots \otimes U_i'^{-1} \otimes \dots \otimes U_m'^{-1}) |\Phi\rangle_u$. Here $U_i'^{-1} \otimes U_i' = I$, and the relation between the results of the measurements and the local unitary operations U_i' is shown in Table III. The notation V_i is the bit value of the Bell-state measurement on the particles x_i and b_{i0} , and P_i is defined as

$$P_i = P_{x_i b_{i0}} \otimes \prod_{k=1}^n P_{b_{ik}}. \quad (19)$$

That is, if $V_i = 0$ and $P_i = +$, Charlie performs the unitary operation U_0 on the i -th photon for reconstructing the unknown state $|\Phi\rangle_u$.

TABLE III: The relation between the values of V_i , P_i and the local unitary operation U_i .

V_i	0	0	1	1
P_i	+	-	+	-
U_i'	U_0	U_1	U_2	U_3

III. DISCUSSION AND SUMMARY

In fact, every one of the multiparty quantum state sharing schemes can be used for multiparty-controlled

teleportation by means that some of the agents act as the controllers. That is, this multiparty QSTS scheme can also be used for completing the task in Ref. [29] efficiently. For sharing of multipartite entanglement, the parties should resort to either multipartite entanglement quantum resources acting as the quantum channel or multipartite entanglement measurements. At present, both of them are not easy to be implemented [35, 36, 37]. With the development of technology, they may be feasible in the future. Compared with the scheme in Ref. [27], this multiparty QSTS scheme is more feasible when the unknown quantum system is m -qubit one and the number of the agents is n ($m, n > 2$) as the sender Alice should perform two $(\frac{n \times m}{2} + 1)$ -photon GHZ-state measurements in the former. Different from the Ref. [29], each of the controllers needs only to take a product measurement $\sigma_x^1 \otimes \sigma_x^2 \otimes \dots \otimes \sigma_x^m$ on his photons, no multipartite entanglement measurements, and the last agent can reconstruct the unknown m -qubit state with only m local unitary operations, not CNOT gates, when he cooperates with the controllers, which makes this QSTS scheme more convenient than that in the former.

From Table III, one can see that the agent Charlie can reconstruct the original unknown state with the probability 100% in principle if he cooperates with all the other agents. However, he will only have the probability $\frac{1}{2^m}$ to get the correct result if one of the other agents does not agree to cooperate as Charlie has only half of the chance to choose the correct operation for each qubit according to the information published by Alice and the other $n-1$ agents. That is, this multiparty QSTS scheme is a (n, n) threshold one, same as that in Ref. [7]. On the other hand, as almost all the quantum source (except for the instances chosen for eavesdropping check) can be used to carry the quantum information if the agents act in concert, the intrinsic efficiency for qubits η_q in this scheme approaches 100%, as same as those in the schemes [28, 29, 34, 38, 39, 40, 41, 42, 43] for quantum teleportation and controlled teleportation. Here [8]

$$\eta_q = \frac{q_u}{q_t}, \quad (20)$$

where q_u is the number of the useful qubits in QSTS and q_t is the number of transmitted qubits. In our scheme for sharing m -qubit quantum information with $n+1$ agents, $q_u = q_t = (n + 1)m$ (Here we exploited the definition of the intrinsic efficiency for qubits introduced by Cabello for quantum cryptography [8]). The total efficiency for this scheme can be calculated as following.

$$\eta_t = \frac{q_u}{q_t + b_t}, \quad (21)$$

where b_t is the number of the classical bits exchanged. $b_t = 2m + nm = (n + 2)m$. That is $\eta_t = \frac{(n+1)m}{(n+1)m + (n+2)m} = \frac{n+1}{2n+3}$, the maximal value for QSTS.

In summary, we present a multiparty QSTS scheme for sharing an arbitrary m -qubit state with m GHZ

states, removing some ideas in the symmetric multiparty-controlled teleportation [29]. It is an optimal one with GHZ states used as quantum channel for sharing an arbitrary m -qubit state as one of the parties needs only to perform two-photon Bell-state measurements and the others use single-photon product measurements. Moreover, its efficiency for qubits is 100%, the maximal value as all the quantum resources are useful in theory, and each of the agents can act as the receiver. Simultaneously, this multiparty QSTS scheme can be used for controlled teleportation efficiently.

ACKNOWLEDGEMENT

This work is supported by the National Natural Science Foundation of China under Grant Nos. 10447106, 10435020, 10254002, A0325401 and 10374010, and Beijing Education Committee under Grant No. XK100270454.

-
- [1] Blakley G R 1979 in *Proceedings of the American Federation of Information Processing 1979 National Computer Conference* (American Federation of Information Processing, Arlington, VA, 1979), p 313
 - [2] Nielsen M A and Chuang I L 2000 *Quantum computation and quantum information* (Cambridge: Cambridge University Press)
 - [3] Gisin N, Ribordy G, Tittel W and Zbinden H 2002 *Rev. Mod. Phys.* **74** 145
 - [4] Long G L and Liu X S 2002 *Phys. Rev. A* **65** 032302
 - [5] Deng F G and Long G L 2003 *Phys. Rev. A* **68** 042315
 - [6] Deng F G and Long G L 2004 *Phys. Rev. A* **70** 012311
 - [7] Hillery M, Bužek V and Berthiaume A 1999 *Phys. Rev. A* **59** 1829
 - [8] Karlsson A and Bourennane M 1998 *Phys. Rev. A* **58** 4394
 - [9] Karlsson A, Koashi M and Imoto N 1999 *Phys. Rev. A* **59** 162
 - [10] Cabello A 2000 *Phys. Rev. Lett.* **85** 5635
 - [11] Deng F G, Zhou H Y and long G L 2005 *Phys. Lett. A* **337** 329
 - [12] Gottesman D 2000 *Phys. Rev. A* **61** 012308
 - [13] Nascimento A C A, Mueller-Quade J and Imai H 2001 *Phys. Rev. A* **64** 042311
 - [14] Yang C P, Gea-Banacloche J 2001 *J. Opt. B: Quantum Semiclass. Opt.* **3** 407
 - [15] Bagherinezhad S and Karimipour V 2003 *Phys. Rev. A* **67**, 044302 (2003).
 - [16] Tyc T and Sanders B C 2002 *Phys. Rev. A* **65** 042310
 - [17] Xiao L, Long G L, Deng F G and Pan J W 2004 *Phys. Rev. A* **69** 052307
 - [18] Deng F G, Long G L and Zhou H Y 2005 *Phys. Lett. A* **340** 43
 - [19] Deng F G et al 2004 *Chin. Phys. Lett.* **21** 2097
 - [20] Guo G P and Guo G C 2003 *Phys. Lett. A* **310** 247
 - [21] Zhang Z J 2005 *Phys. Lett. A* **342** 60
 - [22] Bandyopadhyay S 2000 *Phys. Rev. A* **62** 012308
 - [23] Karimipour V, Bahraminasab A and Bagherinezhad S 2002 *Phys. Rev. A* **65** 042320
 - [24] Zhang Z J, Li Y and Man Z X 2005 *Phys. Rev. A* **71** 044301
 - [25] Deng F G et al 2005 *Phys. Rev. A* **72** 044302
 - [26] Yan F L and Gao T 2005 *Phys. Rev. A* **72** 012304
 - [27] Tittel W, Zbinden H and Gisin N 2001 *Phys. Rev. A* **63** 042301
 - [28] Lance A M, Symul T, Bowen W P, Sanders B C and Lam P K 2004 *Phys. Rev. Lett.* **92** 177903
 - [29] Lance A M, Symul T, Bowen W P, Sanders B C, Tyc T, Ralph T C and Lam P K 2005 *Phys. Rev. A* **71** 033814
 - [30] Cleve R, Gottesman D and Lo H K 1999 *Phys. Rev. Lett.* **83** 648
 - [31] Li Y M, Zhang K S and Peng K C 2004 *Phys. Lett. A* **324** 420
 - [32] Deng F G, Li X H, Li C Y, Zhou P and Zhou H Y 2005 *Phys. Rev. A* **72** 044301
 - [33] Yang C P, Chu S I and Han S Y 2004 *Phys. Rev. A* **70** 022329
 - [34] Deng F G, Li C Y, Li Y S, Zhou H Y and Wang Y 2005 *Phys. Rev. A* **72** 022338
 - [35] Deng F G, Long G L and Liu X S 2003 *Phys. Rev. A* **68** 042317
 - [36] Deng F G and Long G L 2004 *Phys. Rev. A* **69** 052319
 - [37] Deng F G and Long G L 2004 *Preprint* arXiv: quant-ph/0408102
 - [38] Wang C et al 2005 *Phys. Rev. A* **71** 044305
 - [39] Wang C et al 2005 *Opt. Commun.* **253** 15
 - [40] Deutsch D, Ekert A, Jozsa R, Macchiavello C, Popescu S and Sanpera A 1996 *Phys. Rev. Lett.* **77** 2818
 - [41] Murao M, Plenio M B, Popescu S, Vedral V and Knight P L 1998 *Phys. Rev. A* **57** R4075
 - [42] Yang C P and Guo G C 2000 *Chin. Phys. Lett.* **17** 162
 - [43] Rigolin G 2005 *Phys. Rev. A* **71** 032303
 - [44] Deng F G 2005 *Phys. Rev. A* **72** 036301
 - [45] Bouwmeester D, Pan J W, Daniell M, Weinfurter H and Zeilinger A 1999 *Phys. Rev. Lett.* **82** 1345
 - [46] Pan J W, Daniell M, Gasparoni S, Weihs G and Zeilinger A 2001 *Phys. Rev. Lett.* **86**, 4435
 - [47] Zhao Z, Chen Y A, Zhang A N, Yang T, Briegel H J and Pan J W 2004 *Nature* **430** 54
 - [48] Bennett C H, Brassard G, Crépeau C, Jozsa R, Peres A and Wootters W K 1993 *Phys. Rev. Lett.* **70** 1895
 - [49] Gorbachev V N and Trubilko A I 1999 *Preprint* arXiv: quant-ph/9906110.
 - [50] Marinatto L and Weber T 2000 *Found. Phys. Lett.* **13** 119
 - [51] Shi B S, Jiang Y K and Guo G C 2000 *Phys. Lett. A* **268** 161
 - [52] Lu H and Guo G C 2000 *Phys. Lett. A* **276** 209
 - [53] Lee H W 2001 *Phys. Rev. A* **64** 014302
 - [54] Yan F L, Tan H G and Yang L G 2002 *Commun. Theor. Phys.* **37** 649